













Article

Quality Appraisal of Nutritional Guidelines to Prevent, Diagnose, and Treat Malnutrition in All Its Forms during Pregnancy

Cintha Muñoz-Manrique ^{1,2,3} , Mónica Ancira-Moreno ^{1,2,*} , Soraya Burrola-Méndez ^{1,2} , Isabel Omaña-Guzmán ^{1,2} , Elizabeth Hoyos-Loya ¹, Sonia Hernández-Cordero ^{2,4}, Alejandra Trejo-Domínguez ^{1,2} , Mónica Mazariegos ^{2,5} , Natalia Smith ⁶ , Scarlett Alonso-Carmona ^{1,2} , Jennifer Mier-Cabrera ^{1,2} , Loredana Tavano-Colaizzi ^{1,2}, Belén Sánchez-Múzquiz ^{1,2}, Fermín Avendaño-Álvarez ^{1,2}, Karla Muciño-Sandoval ^{1,2}, Nadia C. Rodríguez-Moguel ^{1,2,7} , Magali Padilla-Camacho ^{2,8} , Salvador Espino-y-Sosa ^{2,3} , Lizeth Ibarra-González ¹ and Cristina Medina-Avilés ^{1,2}

- ¹ Health Department, Universidad Iberoamericana, Prolongación Paseo de Reforma 880, Lomas de Santa Fe, Mexico City 01219, Mexico
 - ² Maternal and Child Health and Nutrition Network (MaChiNNe), Observatorio Materno Infantil (OMI), Universidad Iberoamericana, Prolongación Paseo de Reforma 880, Lomas de Santa Fe, Mexico City 01219, Mexico
 - ³ Instituto Nacional de Perinatología, Montes Urales 800 Col., Virreyes Del., Miguel Hidalgo, Mexico City 11000, Mexico
 - ⁴ Research Center for Equitable Development EQUIDE, Universidad Iberoamericana, Prolongación Paseo de Reforma 880, Lomas de Santa Fe, Mexico City 01219, Mexico
 - ⁵ Research Center for the Prevention of Chronic Diseases (CIIPEC), Institute of Nutrition of Central America and Panama (INCAP), Calzada Roosevelt, 6.25 Zona 11, Guatemala City 01011, Guatemala
 - ⁶ Department of Pediatrics, University of California, San Francisco, San Francisco, CA 94143, USA
 - ⁷ Departamento de Investigación en Enfermedades Infecciosas, Instituto Nacional de Enfermedades Respiratorias, Calzada de Tlalpan 4502, Belisario Domínguez Secc 16, Tlalpan, Mexico City 14080, Mexico
 - ⁸ Banco de Leche Humana del Estado de Guanajuato, Vialidad Interior Malvas 367, Revolución, Irapuato 36546, Mexico
- * Correspondence: monica.ancira@ibero.mx; Tel.: +1-360-814-3819



Citation: Muñoz-Manrique, C.; Ancira-Moreno, M.; Burrola-Méndez, S.; Omaña-Guzmán, I.; Hoyos-Loya, E.; Hernández-Cordero, S.; Trejo-Domínguez, A.; Mazariegos, M.; Smith, N.; Alonso-Carmona, S.; et al. Quality Appraisal of Nutritional Guidelines to Prevent, Diagnose, and Treat Malnutrition in All Its Forms during Pregnancy. *Nutrients* **2022**, *14*, 4579. <https://doi.org/10.3390/nu14214579>

Academic Editor: Marloes Dekker Nitert

Received: 26 September 2022

Accepted: 25 October 2022

Published: 1 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: This work aimed to identify clinical practice guidelines (CPGs) that include recommendations for the prevention, diagnosis, and treatment of women's malnutrition during pregnancy and to evaluate the quality of these guidelines using the Appraisal of Guidelines for Research and Evaluation (AGREE II) instrument. We conducted a literature review using PubMed and different websites from January 2009 to February 2021. The quality of the CPGs was independently assessed by reviewers using the AGREE II instrument, which defines guidelines scoring >70% in the overall assessment as "high quality". The analysis included 43 guidelines. Among the main findings, we identified that only half of the CPGs (51.1%) obtained a final "high quality" evaluation. AGREE II results varied widely across domains and categories. The two domains that obtained the highest scores were scope and purpose with 88.3% (range 39 to 100%) and clarity of presentation with 87.2% (range 25 to 100%). Among the "high quality" CPGs, the best scores were achieved by the three guidelines published by the National Institute of Health and Care Excellence (NICE) and the World Health Organization (WHO). Due to the importance of maternal nutrition in pregnancy, it is essential to join forces to improve the quality of the guidelines, especially in CPGs that do not meet the reference standards for quality.

Keywords: AGREE II; clinical practice guidelines; pregnancy; healthy pregnancy; malnutrition

1. Introduction

Maternal nutrition crucially contributes to the offspring's health and future generations. Studies in animals and humans have observed adverse developmental processes

around conception in response to various environmental situations. The quality of the maternal diet, undernutrition, overweight, obesity, and physiological aspects related to glucose and lipid metabolism alteration may affect the embryo's potential with consequences for risk of diseases in the short and long term [1].

Prepregnancy obesity is related to several adverse perinatal outcomes, with the risks increasing as the body mass index increases [2]. Regarding the newborn's nutritional status, due to a proinflammatory status, metabolism alterations in preconception, and other factors, a woman with obesity may have a baby who is large for gestational age. These preconception conditions may also influence placental function and increase the risk of preeclampsia, intrauterine growth restriction, and gestational diabetes mellitus (GDM) [3]. Likewise, women with prepregnancy obesity have a higher probability of excessive gestational weight gain (GWG) and postpartum weight retention [4].

In the same way, the effect of maternal malnutrition is well known. Even though there is a smaller proportion of underweight women worldwide, these proportions are higher in low- and middle-income countries. Women with prepregnancy underweight have a greater probability of having a baby with inadequate fetal growth. These phenotypes may have a detrimental impact on children in their forthcoming years of schooling, educational performance, and health [5].

Women with micronutrient deficiencies before pregnancy, such as iron, folate, iodine, zinc, selenium, and vitamin B complex, are at a greater risk of having these deficiencies during pregnancy as their micronutrient requirements increase to promote adequate fetal development and growth. Deficiencies in critical micronutrients throughout the periconceptional period may affect maternal and offspring health, resulting in early reproductive failure, poor placental vascularization, inadequate fetal brain development, and poor fetal growth. Poor-quality diet intake or altered absorption due to infection, diseases, or inflammation could explain these nutrient deficiencies [6].

Conditions such as inadequate GWG, GDM, and hypertension disorders during pregnancy might influence fetal growth and women's health. Systematic reviews have shown promising results in promoting adequate GWG and glycemic control by following nutrition and lifestyle recommendations during pregnancy [7,8]. In addition, observational studies have reported positive results in preventing hypertensive disorders, low birth weight, and preterm birth in women with better diet quality indices compared to women with lower dietary quality scores [9–11].

The problems of malnutrition mentioned above highlight the need to offer pregnant women food, nutrition, and lifestyle guidelines. International organizations prioritize maternal nutrition in the national public health plan to support a healthy pregnancy. However, women's adherence to nutrition guidelines during pregnancy appears to be below the recommended level [12]. Several sociodemographic factors may influence guideline adherence, although the quality of the recommendations could also affect adherence due to the lack of clear evidence for health professionals to issue these recommendations.

Clinical practice guidelines (CPGs) are proper instruments to implement evidence-based guidelines for patients. However, few studies have evaluated the quality of pregnancy guidelines using a specific methodology. This paper aims to identify CPGs that include recommendations for preventing, diagnosing, and treating women's malnutrition during pregnancy and to evaluate the quality of these guidelines using the Appraisal of Guidelines for Research and Evaluation (AGREE II) instrument.

2. Materials and Methods

2.1. Study Design and Eligibility Criteria

We thoroughly searched CPGs containing nutritional and lifestyle recommendations to prevent, diagnose, and treat all forms of malnutrition during pregnancy. We followed the framework proposed by Arksey and O'Malley [13] and further developed by Levac et al. [14] and the Joanna Briggs Institute [15]. This review process consisted of five stages: (i) identifying the research question; (ii) identifying relevant studies; (iii) study

selection; (iv) charting the data; and (v) collating, summarizing, and reporting results. We added an extra step of (vi) critical appraisal using the AGREE II instrument [16].

The inclusion criteria considered international and national CPGs, consensus, expert opinion based on evidence, and standard references with recommendations on nutritional assessment (behavior, anthropometric, biochemical, clinical evaluation, and lifestyle), healthy diet, dietary modifications, nutritional supplementation, or any nutritional or lifestyle recommendation given in primary healthcare facilities. We excluded opinions, editorials, articles published as communication tools, and CPGs with lifestyle and nutrition recommendations oriented to a specific pathology or associated complications.

2.2. Search Strategy and Studies Selection

We performed a systematic literature search in the database PubMed (<https://ncbi.nlm.nih.gov/pubmed>, accessed on 10 February 2021) and a manual search of guideline-related websites, limiting the search to studies published between January 2009 and February 2021. The selection of this time frame considered the review and update of the Institute of Medicine (IOM) recommendations for weight gain during pregnancy that included ways to encourage their adoption through strategies to assist practitioners with the publication of the “Weight Gain During Pregnancy: Reexamining the Guidelines” in 2009 [17].

For the PubMed search, we used keywords such as pregnancy; pregnancy trimester, first; nutrition assessment; prevention and control; and malnutrition. We followed a Boolean search using search filters to identify any guidelines in English or Spanish (see Table 1). For the guideline-related websites, we selected international agencies and gynecology, obstetrics, and maternal nutrition societies as well as databases from the National Institute of Health and Care Excellence (NICE; UK) and Guidelines International Network (GIN). Key terms of the algorithm were used alone or in combination. We imported all studies identified through the database and the website search to Excel. We checked for and removed duplicates.

Table 1. Boolean search system.

(“Pregnancy”[Mesh] OR “Pregnancy Trimester, First”[Mesh] OR “Pregnancy Trimester, Second”[Mesh] OR “Pregnancy Trimester, Third”[Mesh]) AND (“Prenatal Care”[Mesh] OR “Nutrition Assessment”[Mesh] OR “Nutrition Therapy”[Mesh] OR “prevention and control” [Subheading] OR “Health Promotion”[Mesh]) AND (“Malnutrition”[Mesh] OR “Body Weight”[Mesh] OR “Anemia”[Mesh] OR “Deficiency Diseases”[Mesh] OR “Nutrition Disorders”[Mesh] OR “Gestational Weight Gain”[Mesh] OR “Fetal Growth Retardation”[Mesh] OR “Nutritional Physiological Phenomena”[Mesh])

MeSH: Medical subject headings.

2.3. Quality Assessment

Authors, including dietitians and physicians, participated in the evaluation process. Two authors (CMM and MAM) independently assessed each study’s title and abstract to determine the references’ eligibility. In case of disagreement, another author (SBM) evaluated the guide. Several health professionals (CMA, GAC, JMC, KMS, LIG, LTC, MAM, MPC, NSG, SES, AT, BSM, CMM, FAA, and SHC) conducted a peer review to assess the full text of the potentially eligible documents in order to determine if they meet the inclusion criteria. In case of disagreement, a third author (AT, BSM, CMM, GAC, MPC, and SBM) was responsible to decide whether or not to include it.

We used the AGREE II tool to assess the quality of the selected CPGs. This instrument aims to identify the quality guidelines and their strengths and limitations and consists of 23 key items grouped into six domains: (1) scope and purpose (related to the overall aim of the guideline); (2) stakeholder involvement (measures the extent to which the appropriate stakeholders developed the guideline); (3) rigor of development (concerned with the process used to gather and synthesize the evidence); (4) clarity of presentation (appraises the language, structure, and format of the guideline); (5) applicability (related to the implications of applying the guideline); and (6) editorial independence (evaluates that the formulation of recommendations is unbiased with competing interests) followed by 2 global rating items (“overall assessment”). The evaluation of each item uses a 7-point

Likert rating scale (from 1 = “strongly disagree” to 7 = “strongly agree”), as defined in the AGREE II user’s manual [16].

The overall scores of each domain were calculated by adding their corresponding items and scaling the total as a proportion of the maximum possible score for that domain (max score = 100). The overall assessment requires the user to make a judgment as to the quality of the guideline, taking into account the criteria considered in the assessment process. The user is also asked whether he/she would recommend use of the guideline. A score of >70% indicates “high quality” in the guidelines as determined by the AGREE II user’s manual [16].

Due to time restrictions, only two authors (BSM, LTC, NRM, SES, AT, FAA, JMC, LIG, and MAM) independently evaluated the quality of each CPG using the online AGREE platform, “My AGREE PLUS”, which is suggested by the AGREE II methodology.

2.4. Data Analysis

Means and median scores were calculated in each AGREE II domain to identify the most critical domains across the different guidelines. The overall quality evaluation of each guideline used a threshold of 70% for the final score of each domain. We used Microsoft Excel 2021, version 16.57, for data collection and extraction.

3. Results

Figure 1 provides the PRISMA flow diagram [18] with a detailed summary of the research results. We identified 82 records through the search. After removing duplicated studies, the abstracts of 79 records were screened, leading to 54 CPGs for full review and, finally, 43 for quality assessment.




Figure 1. PRISMA flow diagram of searching and selecting guidelines.

Table 2 shows the general characteristics of the analyzed CPGs. Regarding geographical distribution, 35% of the reviewed guidelines corresponded to the North America Region, with guidelines from Canada (n = 9) and the USA (n = 6), while 37% were from Europe and the Central Asia region, with contribution from the United Kingdom (n = 9), Poland (n = 2), Italy (n = 1), Spain (n = 1), France (n = 1), and Europe region (n = 2). The regions with the lowest number of CPGs reviewed were East Asia and the Pacific, with two CPGs from Australia and the Middle East and North Africa and one CPG from the United Arab Emirates. Finally, we included nine international CPGs (21%).

Table 2. General characteristics of the CPGs included in the analysis.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|--|--|---|---------------------|-------------------|----------------------|--|-------------------------|
| American College of Obstetricians and Gynecologists [19] | ACOG Practice Bulletin No 156: Obesity in Pregnancy | The American College of Obstetricians and Gynecologists | 2015 | USA | 112 | Obstetric care provider or healthcare professionals | P, D |
| American Diabetes Association [20] | Management of diabetes in pregnancy | American Diabetes Association. | 2015 | USA | 23 | Healthcare providers | P, D, T |
| ADA, ASN, Siega-Riz, A.M., King, J.C. [21] | Position of the American Dietetic Association and American Society for Nutrition: obesity, reproduction, and pregnancy outcomes | American Society for Nutrition | 2009 | USA | 97 | Registered dietitians, registered dietetic technicians, and other healthcare professionals | P, D, T |
| Australian Government Department of Health [22] | Clinical Practice Guidelines: Pregnancy Care | Australian Government Department of Health | 2019 | Australia | Not specified | Health professionals | P |
| Bomba-Opoń [23] | Folate supplementation during the preconception period, pregnancy, and puerperium. Polish Society of Gynecologists and Obstetricians Guidelines | Polish Society of Gynecologists and Obstetricians (PTGP): | 2017 | Poland | 30 | Not specified | P, T |
| Cetin, I. [24] | Breastfeeding during pregnancy: position paper of the Italian Society of Perinatal Medicine and the Task Force on Breastfeeding, Ministry of Health, Italy | Italian Society of Perinatal Medicine and Task Force on Breastfeeding | 2014 | Italy | 66 | Not specified | P, T |
| Charzewska, J. [25] | Prophylaxis of vitamin D deficiency, Polish Recommendations 2009 | Not specified | 2010 | Poland | Not specified | Not specified | P, D, T |
| CMACE [26] | Management of Women with Obesity in Pregnancy | Royal College of Obstetricians and Gynecologists | 2010 | UK | 77 | Health professionals | P, D, T |

Table 2. Cont.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|-----------------------|--|--|---------------------|----------------------|----------------------|-----------------------|-------------------------|
| Davies. G.A. [27] | SOGC Clinical Practice Guidelines: Obesity in pregnancy. No. 239, February 2010 | Society of Obstetricians and Gynecologists of Canada | 2010 | Canada | 79 | Obstetrical care team | P, D, T |
| Davies. G.A.L. [28] | No. 239, Obesity in Pregnancy | The Society of Obstetricians and Gynecologists of Canada. | 2018 | Canada | 78 | Obstetrical care team | P, D, T |
| Denison, F.C. [29] | Care of Women with Obesity in Pregnancy: Green-Top Guideline No. 72 | Royal College of Obstetricians and Gynecologists | 2019 | UK | 202 | Healthcare providers | P, D, T |
| Donnay, S. [30] | Iodine supplementation during pregnancy and lactation. Position statement of the working group on disorders related to iodine deficiency and thyroid dysfunction of the Spanish Society of Endocrinology and Nutrition | Spanish Society of Endocrinology and Nutrition | 2014 | Spain | 73 | Not specified | D |
| Fitzsimons, K.J. [31] | Setting maternity care standards for women with obesity in pregnancy | Centre for Maternal and Child Enquiries(CMACE) | 2010 | UK | 55 | Not specified | P, D, T |
| Fleming, N. [32] | Adolescent Pregnancy Guidelines | Canadian Pediatric and Adolescent Gynecology and Obstetricians (CANPAGO) | 2015 | Canada | 119 | Healthcare providers | P, D |
| Haq, A. [33] | Clinical practice guidelines for vitamin D in the United Arab Emirates | Not specified | 2018 | United Arab Emirates | 32 | Physicians | D, T |

Table 2. Cont.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|-------------------|---|---|---------------------|-------------------|----------------------|--------------------------|-------------------------|
| Harden, C.L. [34] | Practice parameter update: management issues for women with epilepsy—focus on pregnancy (an evidence-based review): vitamin K, folic acid, blood levels, and breastfeeding: report of the Quality Standards Subcommittee and Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and American Epilepsy Society | American Academy of Neurology | 2009 | USA | 40 | Not specified | P, T |
| Homer, C.S. [35] | Updated clinical practice guidelines on pregnancy care | Australian Health Ministers' Conference and the Community and Disability Services Ministers' Conference | 2018 | Australia | 11 | Health professionals | P, D |
| Jacob, C.M. [36] | Management of prepregnancy, pregnancy, and postpartum obesity from the FIGO Pregnancy and Non-Communicable Diseases Committee: A FIGO (International Federation of Gynecology and Obstetrics) guideline | International Federation of Gynecology and Obstetrics | 2020 | International | 77 | Healthcare practitioners | P, D, T |
| Koren, G. [37] | Cancer chemotherapy and pregnancy | Society of Obstetricians and Gynaecologists of Canada. | 2013 | Canada | 88 | Healthcare practitioners | P, T |
| Lausman, A. [38] | Intrauterine growth restriction: screening, diagnosis, and management | Society of Obstetricians and Gynaecologists of Canada | 2013 | Canada | 55 | Not specified | P, D, T |
| Marsh, J.C. [39] | Guidelines for the diagnosis and management of aplastic anaemia | General Haematology Task Force of the British Committee for Standards in Haematology | 2009 | UK | 174 | Health professionals | D, T |

Table 2. Cont.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|----------------------|--|--|---------------------|-------------------|----------------------|---|-------------------------|
| Maxwell, C. [40] | Guideline No. 391, Pregnancy and Maternal Obesity Part 1: Preconception and Prenatal Care | Society of Obstetricians and Gynaecologists of Canada (SOGC) | 2019 | Canada | 178 | Healthcare practitioners | P, T |
| Maxwell, C. [41] | Guideline No. 392, Pregnancy and Maternal Obesity Part 2: Team Planning for Delivery and Postpartum Care | Society of Obstetricians and Gynaecologists of Canada (SOGC) | 2019 | Canada | 149 | Healthcare practitioners | P, D, T |
| Mottola, M.F. [42] | 2019 Canadian guideline for physical activity throughout pregnancy | Society of Obstetricians and Gynaecologists of Canada's (SOGC) Maternal Fetal Medicine and Guideline Management and Oversight Committees | 2018 | Canada | 40 | Obstetric care providers, policymakers, and fitness professionals | P |
| Muraro, A. [43] | EAACI food allergy and anaphylaxis guidelines. Primary prevention of food allergy | European Academy of Allergy and Clinical Immunology's (EAACI) | 2014 | Europe | 98 | Medical secondary care, primary care, and nursing | P, T |
| NICE [44] | Antenatal care for uncomplicated pregnancies | National Institute for Health and Care Excellence: Guidelines | 2019 | UK | Not specified | Healthcare providers | P, D, T |
| NICE [45] | Diabetes in pregnancy: management from preconception to the postnatal period | National Institute for Health and Care Excellence: Guidelines | 2020 | UK | Not specified | Healthcare providers | P, D, T |
| NICE [46] | Weight management before, during, and after pregnancy | National Institute for Health and Care Excellence: Guidelines | 2010 | UK | 15 | Health professionals | P, D, T |
| Pavord, S. [47] | UK guidelines on the management of iron deficiency in pregnancy | British Committee for Standards in Haematology | 2012 | UK | 81 | Healthcare professionals | P, D, T |
| Piccinini, V.H. [48] | Canadian Adult Obesity Clinical Practice Guidelines: Weight Management Over the Reproductive Years for Adult Women Living with Obesity | The Canadian Association of Bariatric Physicians and Surgeons (CABPS) | 2020 | Canada | 126 | Primary care providers | P, T |

Table 2. Cont.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|--------------------|---|---|---------------------|-------------------|----------------------|--|-------------------------|
| Ryan, K. [49] | Significant haemoglobinopathies: guidelines for screening and diagnosis | British Society for Haematology | 2010 | UK | 19 | Not specified | D |
| Sentilhes, L. [50] | Shoulder dystocia: guidelines for clinical practice from the French College of Gynecologists and Obstetricians (CNGOF) | French College of Gynecologists and Obstetricians (CNGOF) | 2016 | France | 11 | Obstetricians or Health professional | P, T |
| Siu, A.L. [51] | Screening for Iron Deficiency Anemia and Iron Supplementation in Pregnant Women to Improve Maternal Health and Birth Outcomes: U.S. Preventive Services Task Force Recommendation Statement | U.S. Preventive Services Task Force (USPSTF) | 2015 | USA | 27 | Not specified | D, T |
| SMFM [52] | Society for Maternal–Fetal Medicine Consult Series #50: The role of activity restriction in obstetric management: (Replaces Consult Number 33, August 2014) | Society for Maternal–Fetal Medicine | 2020 | USA | 47 | Obstetricians and maternal–fetal medicinesubspecialists | T |
| WHO [53] | Proper maternal nutrition during pregnancy planning and pregnancy: a healthy start in life | Ministry of Health of Latvia and the WHO Regional Office for Europe | 2017 | Europe | 27 | Not specified | P |
| WHO [54] | Guideline: daily iron supplementation in adult women and adolescent girls | WHO Guidelines Approved by the Guidelines Review Committee | 2016 | International | 28 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |

Table 2. Cont.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|-----------|--|--|---------------------|-------------------|----------------------|--|-------------------------|
| WHO [55] | Guideline: intermittent iron and folic acid supplementation in nonanaemic pregnant women | WHO Guidelines Approved by the Guidelines Review Committee | 2011 | International | 32 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |
| WHO [56] | Guideline: sodium intake for adults and children | WHO Guidelines Approved by the Guidelines Review Committee | 2012 | International | 62 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |
| WHO [57] | Guideline: sugar intake for adults and children, World Health Organization | WHO Guidelines Approved by the Guidelines Review Committee | 2015 | International | 61 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |
| WHO [58] | Guideline: vitamin A supplementation in pregnant women | WHO Guidelines Approved by the Guidelines Review Committee | 2011 | International | 33 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |

Table 2. Cont.

| Reference | Clinical Guideline | Supporting Organization | Year of Publication | Region or Country | Number of References | Target Audience | Type of Recommendations |
|-----------|---|--|---------------------|-------------------|----------------------|--|-------------------------|
| WHO [59] | WHO antenatal care recommendations for a positive pregnancy experience: nutritional interventions update: multiple micronutrient supplements during pregnancy | WHO Guidelines Approved by the Guidelines Review Committee | 2020 | International | 50 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |
| WHO [60] | WHO antenatal care recommendations for a positive pregnancy experience: nutritional interventions update: vitamin D supplements during pregnancy | WHO Guidelines Approved by the Guidelines Review Committee | 2020 | International | 56 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P |
| WHO [61] | WHO recommendations on antenatal care for a positive pregnancy experience | WHO Guidelines Approved by the Guidelines Review Committee | 2016 | International | 214 | Policymakers, expert advisers, and organizations involved in nutrition actions for public health | P, D |

ADA: American Dietetic Association; ASN: American Society of Nutrition; CMACE: Centre for Maternal and Child Enquiries; SMFM: Society for Maternal–Fetal Medicine; NICE: National Institute of Health and Care Excellence; UK: United Kingdom; USA: United States of America; WHO: World Health Organization Type of recommendations—P: prevention; D: diagnosis; T: treatment.

Concerning the type of recommendations in the reviewed CPGs, 86% had recommendations related to prevention, 56% corresponded to diagnosis, and 27% related to treatment (Table 2) of any form of malnutrition during pregnancy.

3.1. Quality of Guidelines According to the AGREE II Domains

Table 3 reports the total score (mean and median) for each domain and the final quality evaluation of all CPGs. Only 51.1% of CPGs obtained a final evaluation of “high quality” with four or more domains reaching a score higher than 70%. Among the “high quality” CPGs, “Antenatal care for uncomplicated pregnancies” (NICE, 2019) [44] had the best score with >90% in all domains, while “Guideline: daily iron supplementation in adult women and adolescent girls” (World Health Organization (WHO), 2016) [54] and “Guideline: vitamin A supplementation in pregnant women” (WHO, 2011) [58] both scored >80% in all six domains.

On the other hand, a CPG published by Bomba-Opoń (2017) [23] and another one published by the American Dietetic Association (ADA), American Society of Nutrition (ASN), Siega-Riz, A.M., King, J.C. (2019) [21] obtained the lowest scores with 17 and 25% of total average, respectively.

When looking more thoroughly at domain scores, the highest domain scores were for “scope and purpose” (domain 1) with a mean score of 88.3% (range 39 to 100%) and “clarity of presentation” (domain 4) with a mean score of 87.2% (range 25 to 100%). Meanwhile, the lowest values were for “applicability” (domain 5) with a mean score of 67.8% (range 4 to 100%) and “rigor and development” (domain 3) with a mean score of 74.3% (range 10 to 100%).

Figures 2 and 3 show the mean and median quality scores of each domain in the guidelines graded as high and low quality.




Figure 2. Average quality score of each domain for all guidelines sorted by quality classification.




Figure 3. Median quality score of each domain for all guidelines sorted by quality classification.

Table 3. Appraisal of Guidelines for Research and Evaluation (AGREE) II version result for clinical practice guidelines.

| Clinical Guideline | AGREE II Domains (%) | | | | | | | Quality of Guidelines |
|---|----------------------|-------------------------|-----------------------|-------------------------|---------------|------------------------|--------------------|-----------------------|
| | Scope and Purpose | Stakeholder Involvement | Rigour of Development | Clarity of Presentation | Applicability | Editorial Independence | Overall Assessment | |
| 12. Management of diabetes in pregnancy [20] | 89 | 92 | 90 | 94 | 88 | 96 | 83 | High quality |
| 2019 Canadian guideline for physical activity throughout pregnancy [42] | 100 | 97 | 92 | 100 | 96 | 88 | 92 | High quality |
| ACOG Practice Bulletin No 156: Obesity in Pregnancy [19] | 83 | 75 | 71 | 94 | 75 | 54 | 67 | Low quality |
| Adolescent Pregnancy Guidelines [32] | 97 | 86 | 81 | 100 | 67 | 75 | 83 | High quality |
| Antenatal care for uncomplicated pregnancies [44] | 100 | 97 | 100 | 100 | 94 | 92 | 100 | High quality |
| Breastfeeding during pregnancy: position paper of the Italian Society of Perinatal Medicine and the Task Force on Breastfeeding, Ministry of Health, Italy [24] | 83 | 69 | 57 | 25 | 38 | 88 | 58 | Low quality |
| Canadian Adult Obesity Clinical Practice Guidelines: Weight Management Over the Reproductive Years for Adult Women Living with Obesity [48] | 94 | 86 | 75 | 92 | 88 | 58 | 83 | High quality |
| Cancer chemotherapy and pregnancy [37] | 89 | 92 | 90 | 94 | 88 | 96 | 83 | High quality |
| Care of Women with Obesity in Pregnancy: Green-top Guideline No. 72 [29] | 86 | 58 | 97 | 97 | 65 | 96 | 83 | High quality |
| Clinical practice guidelines for vitamin D in the United Arab Emirates [33] | 94 | 72 | 51 | 89 | 73 | 29 | 67 | Low quality |
| Clinical Practice Guidelines: Pregnancy Care [22] | 69 | 56 | 56 | 83 | 48 | 71 | 58 | Low quality |
| Diabetes in pregnancy: management from preconception to the postnatal period [45] | 100 | 100 | 84 | 100 | 98 | 92 | 92 | High quality |
| EAACI food allergy and anaphylaxis guidelines. Primary prevention of food allergy [43] | 75 | 89 | 97 | 97 | 85 | 96 | 92 | High quality |
| Folate supplementation during the preconception period, pregnancy and puerperium. Polish Society of Gynecologists and Obstetricians Guidelines [23] | 39 | 28 | 21 | 64 | 4 | 8 | 17 | Low quality |
| Guideline No. 391, Pregnancy and Maternal Obesity Part 1: Preconception and Prenatal Care [40] | 97 | 83 | 100 | 94 | 60 | 83 | 83 | High quality |
| Guideline No. 392, Pregnancy and Maternal Obesity Part 2: Team Planning for Delivery and Postpartum Care [41] | 100 | 100 | 85 | 100 | 96 | 92 | 92 | High quality |
| Guideline: daily iron supplementation in adult women and adolescent girls [54] | 100 | 86 | 98 | 100 | 90 | 100 | 100 | High quality |

Table 3. Cont.

| Clinical Guideline | AGREE II Domains (%) | | | | | | | Quality of Guidelines |
|--|----------------------|-------------------------|-----------------------|-------------------------|---------------|------------------------|--------------------|-----------------------|
| | Scope and Purpose | Stakeholder Involvement | Rigour of Development | Clarity of Presentation | Applicability | Editorial Independence | Overall Assessment | |
| Guideline: intermittent iron and folic acid supplementation in nonanemic pregnant women [55] | 100 | 83 | 96 | 92 | 100 | 83 | 92 | High quality |
| Guideline: sodium intake for adults and children [56] | 100 | 83 | 98 | 100 | 88 | 100 | 92 | High quality |
| Guideline: sugars intake for adults and children. World Health Organization [57] | 100 | 83 | 98 | 100 | 100 | 100 | 92 | High quality |
| Guideline: vitamin A supplementation in pregnant women [58] | 100 | 89 | 98 | 100 | 96 | 100 | 100 | High quality |
| Guidelines for the diagnosis and management of aplastic anemia [39] | 92 | 81 | 65 | 92 | 50 | 58 | 58 | Low quality |
| Intrauterine growth restriction: screening, diagnosis, and management [38] | 81 | 33 | 53 | 89 | 33 | 4 | 50 | Low quality |
| Iodine supplementation during pregnancy and lactation. Position statement of the working group on disorders related to iodine deficiency and thyroid dysfunction of the Spanish Society of Endocrinology and Nutrition [30] | 83 | 56 | 42 | 42 | 27 | 54 | 50 | Low quality |
| Management of prepregnancy, pregnancy, and postpartum obesity from the FIGO Pregnancy and Non-Communicable Diseases Committee: A FIGO (International Federation of Gynecology and Obstetrics) guideline [36] | 100 | 94 | 86 | 100 | 92 | 100 | 92 | High quality |
| Management of Women with Obesity in Pregnancy [26] | 100 | 89 | 92 | 94 | 83 | 92 | 67 | Low quality |
| No. 239-Obesity in Pregnancy [28] | 86 | 39 | 63 | 86 | 33 | 21 | 50 | Low quality |
| Position of the American Dietetic Association and American Society for Nutrition: obesity, reproduction, and pregnancy outcomes [21] | 75 | 47 | 22 | 56 | 25 | 29 | 25 | Low quality |
| Practice parameter update: management issues for women with epilepsy—focus on pregnancy (an evidence-based review): vitamin K, folic acid, blood levels, and breastfeeding: report of the Quality Standards Subcommittee and Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and American Epilepsy Society [34] | 100 | 81 | 73 | 92 | 75 | 75 | 58 | Low quality |
| Proper maternal nutrition during pregnancy planning and pregnancy: a healthy start in life [53] | 89 | 81 | 63 | 83 | 63 | 96 | 67 | Low quality |
| Prophylaxis of vitamin D deficiency: Polish Recommendations 2009 [25] | 58 | 53 | 10 | 81 | 38 | 29 | 50 | Low quality |

Table 3. Cont.

| Clinical Guideline | AGREE II Domains (%) | | | | | | | Quality of Guidelines |
|--|-------------------------|-------------------------|-------------------------|-------------------------|------------------------|------------------------|-------------------------|-----------------------|
| | Scope and Purpose | Stakeholder Involvement | Rigour of Development | Clarity of Presentation | Applicability | Editorial Independence | Overall Assessment | |
| Screening for Iron Deficiency Anemia and Iron Supplementation in Pregnant Women to Improve Maternal Health and Birth Outcomes: U.S. Preventive Services Task Force Recommendation Statement [51] | 97 | 47 | 58 | 69 | 54 | 100 | 58 | Low quality |
| Setting maternity care standards for women with obesity in pregnancy [31] | 83 | 72 | 73 | 83 | 52 | 92 | 67 | Low quality |
| Shoulder dystocia: guidelines for clinical practice from the French College of Gynecologists and Obstetricians (CNGOF) [50] | 81 | 67 | 73 | 78 | 67 | 96 | 67 | Low quality |
| Significant haemoglobinopathies: guidelines for screening and diagnosis [49] | 81 | 92 | 58 | 89 | 58 | 8 | 50 | Low quality |
| Society for Maternal–Fetal Medicine Consult Series #50: The role of activity restriction in obstetric management: (Replaces Consult Number 33, August 2014) [52] | 75 | 44 | 53 | 72 | 42 | 50 | 58 | Low quality |
| SOGC Clinical Practice Guidelines: Obesity in Pregnancy. No. 239, February 2010 [27] | 83 | 39 | 63 | 81 | 31 | 38 | 50 | Low quality |
| UK guidelines on the management of iron deficiency in pregnancy [47] | 75 | 64 | 67 | 78 | 54 | 100 | 67 | Low quality |
| Updated clinical practice guidelines on pregnancy care [35] | 97 | 94 | 90 | 100 | 63 | 88 | 83 | High quality |
| Weight management before, during and after pregnancy [46] | 100 | 100 | 92 | 100 | 94 | 100 | 92 | High quality |
| WHO antenatal care recommendations for a positive pregnancy experience: nutritional interventions update: multiple micronutrient supplements during pregnancy [59] | 89 | 86 | 90 | 86 | 75 | 100 | 83 | High quality |
| WHO antenatal care recommendations for a positive pregnancy experience: nutritional interventions update: vitamin D supplements during pregnancy [60] | 89 | 89 | 90 | 92 | 85 | 100 | 92 | High quality |
| WHO recommendations on antenatal care for a positive pregnancy experience [61] | 89 | 92 | 88 | 92 | 85 | 100 | 92 | High quality |
| Mean (range) | 88.3 (range 39 to 100%) | 75.4 (range 28 to 100%) | 74.3 (range 10 to 100%) | 87.2 (range 25 to 100%) | 67.8 (range 4 to 100%) | 75 (range 4 to 100%) | 72.9 (range 17 to 100%) | |
| Median (range) | 89 (range 39 to 100%) | 83 (range 28 to 100%) | 81 (range 10 to 100%) | 92 (range 25 to 100%) | 73 (range 4 to 100%) | 92 (range 4 to 100%) | 83 (range 17 to 100%) | |

EAACI: European Academy of Allergy and Clinical Immunology; WHO: World Health Organization.

3.1.1. Scope and Purpose Domain

Domain 1 had the highest results with a mean of 88.3% (range 39 to 100%) and a median of 89% (range 39 to 100%). Moreover, 30.2% (n = 13) of the guidelines had a maximum score of 100%. Only one guideline presented a score below 50%, which was the Bomba-Opoń (2017) guideline [23] with 39%.

3.1.2. Stakeholder Involvement

The mean score in domain 2 was 75.4% (range 28 to 100%), and the median was 83% (range 28 to 100%). Only 67.4% (n = 29) of the guidelines received scores higher than 70%. The guideline published by Maxwell, C. (2017) [41] and those by NICE in 2010 [45] and 2020 [46] had the highest scores (100% each). The Bomba-Opoń guideline [23] scored the lowest score of 28%.

3.1.3. Rigor of Development

Regarding domain 3, the best score was 100% (range 10 to 100%), which was obtained by Maxwell, C. (2019) [40] and NICE (2019) [44]. In contrast, the lowest scores were 21 and 22% for the guidelines by ADA, ASN, Siega-Riz, A.M., and King, J.C. [21] and Bomba-Opoń [23], respectively.

3.1.4. Clarity of Presentation

The mean score in domain 4 was the second highest with 87.2% (range 25 to 100%). Only 27.9% (n = 12) of the guidelines had a high score (100%). In contrast, the guidelines of Cetin, I. [24] and Donnay, S. [30] obtained the lowest scores of 25 and 42%, respectively.

3.1.5. Applicability

Domain 5 had the worst result, with the lowest mean score of 67.8% (range 4 to 100%) and a median of 73% (range 4 to 100%). Only 51.1% (n = 22) of the reviewed guidelines obtained a score >70%, and 25.5% (n = 11) had a score below or equal to 50%. The WHO guidelines [55,57] received a high score of 100%. Once again, the guidelines by Bomba-Opoń [23] and ADA, ASN, Siega-Riz, A.M., and King, J.C. [21] obtained the lowest scores with 4 and 25%, respectively.

3.1.6. Editorial Independence

For domain 6, the mean score was 75% (range 4 to 100%), and the median score was 92% (range 4 to 100%). Nine of the CPGs analyzed (20.9%) received a scores below or equal to 50%. Compared with the others, this domain obtained the lowest scores (e.g., Lausman, A. [38] obtained 4%, Ryan, K. [49] obtained 8%, and [23] also obtained 8%).

4. Discussion

The current study presents a rigorous review that explores the quality of 43 CPGs designed to prevent, diagnose, and treat malnutrition in pregnancy using the AGREE II tool. Among the main findings, we identified that only half of the CPGs (51.1%) obtained a final “high quality” evaluation with four or more domains reaching a score higher than 70%. The guidelines classified as high and low quality had a higher evaluation in the classifications of domain 1 “scope and purpose” (mean = 88.3%, range = 39–100%) and domain 4 “clarity of presentation” (mean = 87.2%, range = 25–100%). The domains with the lowest score in both groups were “applicability” (mean = 67.8%, range = 4–100%), “rigor of development” (mean = 74.3.3%, range = 10–100%), and “editorial independence” (mean = 75.0%, range = 4–100%).

To our knowledge, there are few systematic reviews that have evaluated the quality of nutrition care guidelines in this population [62–64]. These systematic reviews focused mainly on the assessment of weight management guidelines. The findings published by all of them are consistent with ours. The quality of the guidelines regarding the “rigor of development” and “applicability” domains is poor. These observations impact the

failure to establish recommendations in clinical practice and, consequently, to address malnutrition in all its forms in pregnant women. As international organizations recognize them, all the guidelines identified the objective and purpose of care in clinical practice. However, it is necessary to go beyond the purpose because clinical practice guidelines are developed to improve the medical decision-making process among health professionals [65]. Likewise, considering other nutrition indicators, such as food intake, diet quality, and nutrition biomarkers, to assess clinical nutrition guidelines may improve patient-centered antenatal care.

Pregnancy is a unique time for identifying, preventing, and correcting any form of malnutrition in women, which may impact the health of the next generation. This opportunity is of utmost importance as this is a sensitive time, and most countries integrate antenatal care into universal health access. Dietary interventions to promote a healthy diet during pregnancy are part of the double-duty actions suggested by a group of experts to end malnutrition in all its form by 2030 [66]. However, from a nutrition care approach, it is recommended that the nutrition care process be followed to establish specific and evidence-based dietary interventions for each person. In this context, researchers and professionals should strengthen the nutrition methodology in research studies and consider the best evidence-based clinical nutrition practice guidelines.

Assessment of the quality of CPGs is relevant in the clinical field and for formulating and implementing programs and interventions to improve the quality of care and the population's nutritional status. We recommend that guideline developers, clinicians, researchers, and policymakers consider and use the AGREE II instrument as it is detailed and easy to apply in situations of poor nutrition in pregnant women and other clinical conditions [64]. In addition, this tool allows critical evaluation before implementation, which would support decision-making around the health system of a country or region and even during emergencies.

Maternal nutritional status during pregnancy plays a critical role in fetal growth and development, contributing to the offspring's health and future maternal health; therefore, improving the quality of the domains with lower scores is essential. Enhancing the quality of CPGs could improve nutritional care during this life stage and the health of mothers and their offspring across generations. Due to the importance of maternal nutrition in pregnancy, it is essential to join forces to improve the quality of the guidelines, especially CPGs that need to meet the reference standards for quality.

Author Contributions: Conceptualization, M.A.-M., S.B.-M., and C.M.-M.; methodology, N.S., S.B.-M., M.A.-M., and I.O.-G.; formal analysis, S.H.-C., A.T.-D., and I.O.-G.; investigation, C.M.-M., M.M., N.S., L.T.-C., F.A.-Á., S.E.-y.-S., K.M.-S., L.I.-G., J.M.-C., S.A.-C., B.S.-M., N.C.R.-M., M.P.-C., and C.M.-A.; resources, S.B.-M., M.A.-M., and S.H.-C.; data curation, M.A.-M., N.S., and S.B.-M.; writing—original draft preparation, S.B.-M., M.A.-M., C.M.-M., I.O.-G., and E.H.-L.; writing—review and editing, S.B.-M., M.A.-M., C.M.-M., I.O.-G., E.H.-L., J.M.-C., and S.H.-C.; supervision, M.A.-M., S.B.-M., N.S., I.O.-G., and S.H.-C.; project administration, M.A.-M., and I.O.-G.; funding acquisition, M.A.-M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the United Nations Children's Fund (UNICEF; RFP9156422).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Fleming, T.P.; Watkins, A.J.; Velazquez, M.A.; Mathers, J.C.; Prentice, A.M.; Stephenson, J.; Barker, M.; Saffery, R.; Yajnik, C.S.; Eckert, J.J.; et al. Origins of Lifetime Health Around the Time of Conception: Causes and Consequences. *Lancet* **2018**, *391*, 1842–1852. [\[CrossRef\]](#)
2. Santos, S.; Voerman, E.; Amiano, P.; Barros, H.; Beilin, L.J.; Bergström, A.; Charles, M.A.; Chatzi, L.; Chevrier, C.; Chrousos, G.P.; et al. Impact of maternal body mass index and gestational weight gain on pregnancy complications: An individual participant data meta-analysis of European, North American and Australian cohorts. *BJOG Int. J. Obs. Gynaecol.* **2019**, *126*, 984–995. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Poston, L.; Caleyachetty, R.; Cnattingius, S.; Corvalán, C.; Uauy, R.; Herring, S.; Gillman, M. Preconceptional and maternal obesity: Epidemiology and health consequences. *Lancet Diabetes Endocrinol.* **2016**, *4*, 1025–1036. [\[CrossRef\]](#)
4. Nehring, I.; Schmoll, S.; Beyerlein, A.; Hauner, H.; Von Kries, R. Gestational weight gain and long-term postpartum weight retention. *Am. J. Clin. Nutr.* **2011**, *94*, 1225–1231. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Victora, C.G.; Adair, L.; Fall, C.; Hallal, P.C.; Martorell, R.; Richter, L.; Singh, S.H.; Maternal and Child Undernutrition Study Group. Maternal and child undernutrition: Consequences for adult health and human capital. *Lancet* **2008**, *371*, 340–357. [\[CrossRef\]](#)
6. Bailey, R.L.; West, K.P.; Black, R.E. The epidemiology of global micronutrient deficiencies. *Ann. Nutr. Metab.* **2015**, *66*, 22–33. [\[CrossRef\]](#)
7. International Weight Management in Pregnancy (i-WIP) Collaborative Group. Effect of diet and physical activity based interventions in pregnancy on gestational weight gain and pregnancy outcomes: Meta-analysis of individual participant data from randomised trials. *BMJ* **2017**, *358*, j3119.
8. Yamamoto, J.M.; Kellett, J.E.; Balsells, M.; García-Patterson, A.; Hadar, E.; Solà, I.; Gich, I.; van der Beek, E.M.; Castañeda-Gutiérrez, E.; Heinonen, S. Gestational diabetes mellitus and diet: A systematic review and meta-analysis of randomized controlled trials examining the impact of modified dietary interventions on maternal glucose control and neonatal birth weight. *Diabetes Care* **2018**, *41*, 1346–1361. [\[CrossRef\]](#)
9. Ancira-Moreno, M.; O'Neill, M.S.; Rivera-Dommarco, J.Á.; Batis, C.; Rodríguez-Ramírez, S.; Sánchez, B.N.; Castillo-Castrejon, M.; Vadillo-Ortega, F. Dietary patterns and diet quality during pregnancy and low birthweight: The PRINCESA cohort. *Matern. Child Nutr.* **2020**, *16*, e12972. [\[CrossRef\]](#)
10. Gete, D.G.; Waller, M.; Mishra, G.D. Effects of maternal diets on preterm birth and low birth weight: A systematic review. *Br. J. Nutr.* **2020**, *123*, 446–461. [\[CrossRef\]](#)
11. Li, M.; Grewal, J.; Hinkle, S.N.; Yisahak, S.F.; Grobman, W.A.; Newman, R.B.; Skupski, D.W.; Chien, E.K.; Wing, D.A.; Grantz, K.L.; et al. Healthy dietary patterns and common pregnancy complications: A prospective and longitudinal study. *Am. J. Clin. Nutr.* **2021**, *114*, 1229–1237. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Caut, C.; Leach, M.; Steel, A. Dietary guideline adherence during preconception and pregnancy: A systematic review. *Matern. Child Nutr.* **2020**, *16*, e12916. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Arksey, H.; O'Malley, L. Scoping studies: Towards a methodological framework. *Int. J. Soc. Res. Methodol. Theory Pract.* **2005**, *8*, 19–32. [\[CrossRef\]](#)
14. Levac, D.; Colquhoun, H.; O'Brien, K.K. Scoping studies: Advancing the methodology. *Implement. Sci.* **2010**, *5*, 69. [\[CrossRef\]](#) [\[PubMed\]](#)
15. The Joanna Briggs Institute. *Joanna Briggs Institute Reviewer's Manual: 2015 Edition/Supplement*; The Joanna Briggs Institute: Adelaide, Australia, 2015; pp. 6–23.
16. Brouwers, M.C.; Kho, M.E.; Browman, G.P.; Burgers, J.S.; Cluzeau, F.; Feder, G.; Fervers, B.; Graham, I.D.; Grimshaw, J.; Hanna, S.E.; et al. AGREE II: Advancing guideline development, reporting and evaluation in healthcare. *Prev. Med.* **2010**, *51*, 421–424. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines. *Weight Gain During Pregnancy: Reexamining the Guidelines*; Rasmussen, K.M., Yaktine, A.L., Eds.; National Academies Press US: Washington, DC, USA, 2009.
18. Page, M.J.; Moher, D.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. PRISMA 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews. *BMJ* **2021**, *372*, n160. [\[CrossRef\]](#)
19. ACOG Practice Bulletin No 156: Obesity in Pregnancy. *Obs. Gynecol.* **2015**, *126*, e112–e126. [\[CrossRef\]](#)
20. American Diabetes Association. 12. Management of diabetes in pregnancy. *Diabetes Care.* **2015**, *38*, S77–S79. [\[CrossRef\]](#)
21. Siega-Riz, A.M.; King, J.C. Position of the American Dietetic Association and American Society for Nutrition: Obesity, reproduction, and pregnancy outcomes. *J. Am. Diet. Assoc.* **2009**, *109*, 918–927.
22. Department of Health. *Clinical Practice Guidelines: Pregnancy Care*; Australian Government Department of Health: Canberra, Australia, 2018; pp. 9–302.
23. Bomba, O.D.; Hirnle, L.; Kalinka, J.; Seremak, M.A. Folate supplementation during the preconception period, pregnancy and puerperium. Polish Society of Gynecologists and Obstetricians Guidelines. *Ginekol. Polska.* **2017**, *88*, 633–636. [\[CrossRef\]](#)

24. Cetin, I.; Assandro, P.; Massari, M.; Sagone, A.; Gennaretti, R.; Donzelli, G.; Knowles, A.; Monasta, L.; Davanzo, R.; Working Group on Breastfeeding; et al. Breastfeeding during pregnancy: Position paper of the Italian Society of Perinatal Medicine and the Task Force on Breastfeeding, Ministry of Health, Italy. *J. Hum. Lact.* **2014**, *30*, 20–27. [\[CrossRef\]](#)
25. Charzewska, J.; Chlebna-Sokół, D.; Chybicka, A.; Czech-Kowalska, J.; Dobrzańska, A.; Helwich, E.; Imiela, J.R.; Karczmarewicz, E.; Ksiazyk, J.B.; Lewiński, A.; et al. Recommendations of prophylaxis of vitamin D deficiency in Poland (2009). *Med. Wieku Rozw.* **2010**, *14*, 218–223.
26. Modder, J.; Fitzsimons, K. Management of women with obesity in pregnancy. In *CMACE/RCOG Joint Guideline*; Public Health Agency: Ottawa, ON, Canada, 2010; pp. 1–29.
27. Davies, G.A.L.; Maxwell, C.; McLeod, L.; Gagnon, R.; Basso, M.; Bos, H.; Delisle, M.F.; Farine, D.; Hudon, L.; Menticoglou, S.; et al. SOGC Clinical Practice Guidelines: Obesity in pregnancy. No. 239, February 2010. *Int. J. Gynaecol. Obstet.* **2010**, *110*, 167–173. [\[CrossRef\]](#)
28. Davies, G.A.L.; Maxwell, C.; McLeod, L. No. 239-Obesity in Pregnancy. *J. Obstet. Gynaecol. Can.* **2018**, *40*, e630–e639. [\[CrossRef\]](#) [\[PubMed\]](#)
29. Denison, F.C.; Aedla, N.R.; Keag, O.; Hor, K.; Reynolds, R.M.; Milne, A.; Diamond, A.; the Royal College of Obstetricians and Gynaecologists. Care of Women with Obesity in Pregnancy. *BJOG* **2018**, *126*, e62–e106. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Donnay, S.; Arena, J.; Lucas, A.; Velasco, I.; Ares, S.; Working Group on Disorders Related to Iodine Deficiency and Thyroid Dysfunction of the Spanish Society of Endocrinology and Nutrition. Iodine supplementation during pregnancy and lactation. Position statement of the working group on disorders related to iodine deficiency and thyroid dysfunction of the Spanish Society of Endocrinology and Nutrition. *Endocrinol. Nutr.* **2014**, *61*, 27–34. [\[CrossRef\]](#)
31. Fitzsimons, K.J.; Modder, J.; Centre for Maternal and Child Enquires. Setting maternity care standards for women with obesity in pregnancy. *Semin. Fetal Neonatal Med.* **2010**, *15*, 100–107. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Fleming, N.; O'Driscoll, T.; Becker, G.; Spitzer, R.F.; Canpago Committee. Adolescent Pregnancy Guidelines. *J. Obstet. Gynaecol. Can.* **2015**, *37*, 740–756. [\[CrossRef\]](#)
33. Haq, A.; Wimalawansa, S.J.; Pludowski, P.; Anouti, F.A. Clinical practice guidelines for vitamin D in the United Arab Emirates. *J. Steroid Biochem. Mol. Biol.* **2018**, *175*, 4–11. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Harden, C.L.; Pennell, P.B.; Koppel, B.S.; Hovinga, C.A.; Gidal, B.; Meador, K.J.; Hopp, J.; Ting, T.Y.; Hauser, W.A.; Thurman, D.; et al. American Epilepsy Society. Practice parameter update: Management issues for women with epilepsy—focus on pregnancy (an evidence-based review): Vitamin K, folic acid, blood levels, and breastfeeding; Report of the Quality Standards Subcommittee and Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and American Epilepsy Society. *Neurology* **2009**, *73*, 142–149.
35. Homer, C.S.; Oats, J.; Middleton, P.; Ramson, J.; Diplock, S. Updated clinical practice guidelines on pregnancy care. *Med. J. Aust.* **2018**, *209*, 409–412. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Jacob, C.M.; Killeen, S.L.; McAuliffe, F.M.; Stephenson, J.; Hod, M.; Diaz-Yamal, I.; Malhotra, J.; Mocanu, E.; McIntyre, H.D.; Kihara, A.B.; et al. Prevention of noncommunicable diseases by interventions in the preconception period: A FIGO position paper for action by healthcare practitioners. *Int. J. Gynecol. Obstet.* **2020**, *151*, 6–15. [\[CrossRef\]](#) [\[PubMed\]](#)
37. Koren, G.; Carey, N.; Gagnon, R.; Maxwell, C.; Nulman, I.; Senikas, V. Cancer chemotherapy and pregnancy. *J. Obstet. Gynaecol. Can.* **2013**, *35*, 263–278. [\[CrossRef\]](#)
38. Lausman, A.; Kingdom, J.; Maternal Fetal Medicine Committee. Intrauterine growth restriction: Screening, diagnosis, and management. *J. Obstet. Gynaecol. Can.* **2013**, *35*, 741–748. [\[CrossRef\]](#)
39. Marsh, J.C.W.; Ball, S.E.; Cavenagh, J.; Darbyshire, P.; Dokal, I.; Gordon-Smith, E.C.; Keidan, J.; Laurie, A.; Martin, A.; Mercieca, J.; et al. Guidelines for the diagnosis and management of aplastic anaemia. *Br. J. Haematol.* **2009**, *147*, 43–70. [\[CrossRef\]](#)
40. Maxwell, C.; Gaudet, L.; Cassir, G.; Nowik, C.; Lynne, M.L.; Jacob, C.E.; Walker, M. Guideline No. 391-Pregnancy and Maternal Obesity Part 1: Pre-conception and Prenatal Care. *J. Obstet. Gynaecol. Can.* **2019**, *41*, 1623–1640. [\[CrossRef\]](#)
41. Maxwell, C.; Gaudet, L.; Cassir, G.; Nowik, C.; Lynne, M.L.; Jacob, C.E.; Walker, M. Guideline No. 392-Pregnancy and Maternal Obesity Part 2: Team Planning for Delivery and Postpartum Care. *J. Obstet. Gynaecol. Can.* **2019**, *41*, 1660–1675. [\[CrossRef\]](#)
42. Mottola, M.F.; Davenport, M.H.; Ruchat, S.M.; Davies, G.A.; Poitras, V.J.; Gray, C.E.; Jaramillo-Garcia, A.; Barrowman, N.; Adamo, K.B.; Duggan, M.; et al. 2019 Canadian guideline for physical activity throughout pregnancy. *Br. J. Sports Med.* **2018**, *52*, 1339–1346. [\[CrossRef\]](#)
43. Muraro, A.; Halken, S.; Arshad, S.H.; Beyer, K.; Dubois, A.E.J.; Du-Toit, G.; Eigenmann, P.A.; Grimshaw, K.E.C.; Hoest, A.; Lack, G.; et al. EAACI food allergy and anaphylaxis guidelines. Primary prevention of food allergy. *Allergy* **2014**, *69*, 590–601. [\[CrossRef\]](#)
44. NICE (National Institute for Health and Care Excellence). *Antenatal Care for Uncomplicated Pregnancies*; NICE: London, UK, 2019; pp. 4–60.
45. NICE (National Institute of Health and Care Excellence). *Diabetes in Pregnancy: Management from Preconception to the Postnatal Period*; NICE: London, UK, 2020; pp. 4–39.
46. NICE (National Institute of Health and Care Excellence). *Weight Management before, during and after Pregnancy*; NICE: London, UK, 2010; pp. 5–52.
47. Pavord, S.; Myers, B.; Robinson, S.; Allard, S.; Strong, J.; Oppenheimer, C.; on behalf of the British Committee for Standards in Haematology. UK guidelines on the management of iron deficiency in pregnancy. *Br. J. Haematol.* **2012**, *156*, 588–600. [\[CrossRef\]](#)

48. Piccinini, V.H.; Adamo, K.; Bell, R.; Pereira, L.; Nerenberg, K. *Canadian Adult Obesity Clinical Practice Guidelines: Weight Management Over the Reproductive Years for Adult Women Living with Obesity*; CMAJ: Kingston, ON, Canada, 2020; pp. 1–12.
49. Ryan, K.; Bain, B.J.; Worthington, D.; James, J.; Plews, D.; Mason, A.; Roper, D.; Rees, D.C.; De la Salle, B.; Streetly, A.; et al. Significant haemoglobinopathies: Guidelines for screening and diagnosis. *Br. J. Haematol.* **2010**, *149*, 35–49. [[CrossRef](#)] [[PubMed](#)]
50. Sentilhes, L.; Sénat, M.V.; Boulogne, A.I.; Deneux-Tharaux, C.; Fuchs, F.; Legendre, G.; Le Ray, C.; Lopez, E.; Schmitz, T.; Lejeune-Saada, V. Shoulder dystocia: Guidelines for clinical practice from the French College of Gynecologists and Obstetricians (CNGOF). *Eur. J. Obstet. Gynecol. Reprod. Biol.* **2016**, *203*, 156–161. [[CrossRef](#)] [[PubMed](#)]
51. Siu, A.L.; U.S. Preventive Services Task Force. Screening for Iron Deficiency Anemia and Iron Supplementation in Pregnant Women to Improve Maternal Health and Birth Outcomes: U.S. Preventive Services Task Force Recommendation Statement. *Ann. Intern. Med.* **2015**, *163*, 529–536. [[CrossRef](#)]
52. Society for Maternal-Fetal Medicine (SMFM); Lauder, J.; Sciscione, A.; Biggio, J.; Osmundson, S. Society for Maternal-Fetal Medicine Consult Series #50: The role of activity restriction in obstetric management: (Replaces Consult Number 33, August 2014). *Am. J. Obstet Gynecol.* **2020**, *223*, B2–B10. [[PubMed](#)]
53. Meija, L.; Rezeberga, D. *Proper Maternal Nutrition during Pregnancy Planning and Pregnancy: A Healthy Start in Life*; WHO: Copenhagen, Dinamarca, 2017; pp. 4–31.
54. World Health Organization. *Guideline: Daily Iron Supplementation in Adult Women and Adolescent Girls*; World Health Organization: Geneva, Switzerland, 2016; pp. 1–25.
55. World Health Organization. *Guideline: Intermittent Iron and Folic Acid Supplementation in Menstruating Women*; World Health Organization: Geneva, Switzerland, 2011; pp. 5–25.
56. World Health Organization. *Guideline: Sodium Intake for Adults and Children*; World Health Organization: Geneva, Switzerland, 2012; pp. 1–42.
57. World Health Organization. *Guideline: Sugars Intake for Adults and Children*; World Health Organization: Geneva, Switzerland, 2015; pp. 1–45.
58. World Health Organization. *Guideline: Vitamin A Supplementation in Pregnant Women*; World Health Organization: Geneva, Switzerland, 2011; pp. 1–25.
59. World Health Organization. *WHO Antenatal Care Recommendations for a Positive Pregnancy Experience: Nutritional Interventions Update: Multiple Micronutrient Supplements during Pregnancy*; World Health Organization: Geneva, Switzerland, 2020; pp. 1–122.
60. World Health Organization. *WHO Antenatal Care Recommendations for a Positive Pregnancy Experience: Nutritional Interventions Update: Vitamin D Supplements during Pregnancy*; World Health Organization: Geneva, Switzerland, 2020; pp. 1–34.
61. World Health Organization. *WHO Recommendations on Antenatal Care for A Positive Pregnancy Experience*; World Health Organization: Geneva, Switzerland, 2016; pp. 1–152.
62. Grammatikopoulou, M.G.; Theodoridis, X.; Gkiouras, K.; Lampropoulou, M.; Petalidou, A.; Patelida, M.; Tsiros, E.; Papoutsakis, C.; Goulis, D.G. Methodological quality of clinical practice guidelines for nutrition and weight gain during pregnancy: A systematic review. *Nutr. Rev.* **2020**, *78*, 546–562. [[CrossRef](#)]
63. Simon, A.; Pratt, M.; Hutton, B.; Skidmore, B.; Fakhraei, R.; Rybak, N.; Corsi, D.J.; Walker, M.; Velez, M.P.; Smith, G.N.; et al. Guidelines for the management of pregnant women with obesity: A systematic review. *Obes. Rev.* **2020**, *21*, e12972. [[CrossRef](#)]
64. Harrison, C.L.; Teede, H.; Khan, N.; Lim, S.; Chauhan, A.; Drakeley, S.; Moran, L.; Boyle, J. Weight management across preconception, pregnancy, and postpartum: A systematic review and quality appraisal of international clinical practice guidelines. *Obes. Rev.* **2021**, *22*, e13310. [[CrossRef](#)]
65. Brouwers, M.C.; Kho, M.E.; Browman, G.P.; Burgers, J.S.; Cluzeau, F.; Feder, G.; Fervers, B.; Graham, I.D.; Hanna, S.E.; Makarski, J.; et al. Development of the AGREE II, part 1: Performance, usefulness and areas for improvement. *CMAJ* **2010**, *182*, 1045–1052. [[CrossRef](#)]
66. Hawkes, C.; Ruel, M.T.; Salm, L.; Sinclair, B.; Branca, F. Double-duty actions: Seizing programme and policy opportunities to address malnutrition in all its forms. *Lancet* **2020**, *395*, 142–155. [[CrossRef](#)]